

Swimming performance and dry-land upper limbs strength in age group swimmers

Nuno Amaro¹, Daniel A. Marinho^{2, 4}, Mário C. Marques^{2, 4}, Nuno Batalha^{3, 4}, Pedro G. Morouço¹

1. Polytechnic Institute of Leiria, Leiria, Portugal; 2. University of Beira Interior, Covilhã, Portugal; 3. Departamento de Desporto e Saúde, Escola de Ciência e Tecnologia, Universidade de Évora, Portugal; 4. Research Centre of Sports, Health and Human Development, CIDESD, STRONG Research Community; pedro.morouco@ipleiria.pt

High values of dry-land strength and power, particularly in the upper-body, have been identified as a determinant factor to success in competitive swimming. Ball throwing is a common test used to assess dry-land strength and power in age group swimmers. Therefore, the aim of this study was to determine the correlation between ball throwing measures and swimming performance. Twenty-one male prepubescent swimmers (12.7 ± 0.7 years; 47.7 ± 9.6 Kg; 1.56 ± 0.78 m) were recruited for this research. Experiments were conducted during the competitive period of the spring training, thus ensuring that the subjects were in a prime training period cycle. After a standard warm-up of articular mobilization and rope skipping of approximately 10 minutes, Ball throwing (BT) distances (in m) were measured through maximal throwing velocity tests using a 1 and 3 kg medicine ball. Preceding the tests, each participant executed several random throws for warm-up, with both balls. Each participant executed three throws with 2 minutes rest between attempts. Three technical valid attempts were used to calculate the average for analysis. Swimming performance tests were executed after an 800 m moderate intensity warm-up in a 25 m indoor swimming pool. All subjects completed one maximal test of 50 m and 15 m (to exclude the influence of start) in front crawl in order to access their best time in each test (t50 and t15). Short distances were chosen due to the influence of force application over these distances. Spearman correlation coefficients were calculated between in water and dry land parameters assessed. Significance was accepted at the $p < 0.05$ level. Correlations obtained between ball throwing values and swimming performance were moderate to strong for both 1 and 3 Kg balls throws (table 1). Mean values (\pm sd) obtained for ball throwing were: 4.19 ± 0.71 m (1 kg) and 2.84 ± 0.47 m (3 kg). Swimming performance mean values (\pm sd) were 10.63 ± 0.51 s for 15 m trial and 33.70 ± 2.46 s for 50 m trial. Results showed moderate to strong associations between variables, which may indicate that ball throwing can be a valid procedure to evaluate age group swimmers; however, it does not clearly explain performance variability. Lack of specificity of this dry-land test in relation to in-water tests can explain the results.

Table 1
Correlation values (ρ) between ball throwing values (1 and 3 kg) and swimming performance (15 and 50 m).

	BT_1kg	BT_3KG
t15	-0.63*	-0.58*
t50	-0.60*	-0.64*

* $p < 0.05$

Can a Halliwick swimming programme develop water competence, static and dynamic balance in disabled participants?

Andreia Maia¹, Jean Amarante¹, Nuno Serra¹, Carolina Vila-Chã^{1,3}, Tiago M. Barbosa^{2,3}, Mário J. Costa^{1,3}

1. Polytechnic Institute of Guarda, Portugal; 2. Nanyang Technological University, Singapore; 3. Research Centre in Sport Sciences, Health Sciences and Human Development, CIDESD, Portugal; mario.costa@ipg.pt

The Halliwick concept is an aquatic rehabilitation program aiming to enhance balance and core stability of disabled individuals. Previous studies focused on assessing the participants' satisfaction on Halliwick programs or the acquisition of water motor skills (Garcia et al., 2012). Yet, as far as our understanding goes there is not in the literature the assessment of the water independence or the transfer to land-based body balance and posture. Nine individuals (33 ± 12.3 years) diagnosed with cerebral palsy ($n=2$), mental disability ($n=4$) and trisomy 21 ($n=3$) were took part in a Halliwick's 10-point programme. The programme had a duration of 15 weeks comprising one weekly session with one hour of duration. All sessions were planned and implemented by a trained therapist. In the beginning (W0) and in the end (W15) of the intervention programme it was assessed the water competence, static and dynamic balance. Water competence was assessed by the Water Orientation Test Align 2 (WOTA, in points) that is capped to 81 points (Tiroshi et al., 2008). Static balance was assessed by one-leg stance test (OST, in s) and functional reaching test (FRT, in m) according to standard guidelines. The Time up & go test (TUGT, in s) was selected as a measure of

dynamic balance. Water competence measured by WOTA 2 showed significant improvements ($WOTA_{W0} = 39 \pm 11$ vs $WOTA_{W15} = 57 \pm 16$ points, $p < 0.01$) between the beginning and end of the programme. Static balance also improved. There was a significant change in the OST ($OST_{W0} = 15.56 \pm 12.88$ vs $OST_{W15} = 17.78 \pm 12.48$ s, $p = 0.04$) and FRT ($FRT_{W0} = 0.20 \pm 0.09$ vs $FRT_{W15} = 0.24 \pm 0.11$ m, $p < 0.01$) performances. Dynamic balance also improved ($TUGT_{W0} = 9.12 \pm 2.61$ vs $TUGT_{W15} = 7.74 \pm 2.52$ s, $p = 0.01$) after the 15th week. It can be concluded that 15 weeks of a well-designed Halliwick programme can improve water competence of disabled individuals. Concurrently, it was also noted a positive transfer of the skills acquired in water to on land body balance and posture. Hence, fitness and health practitioners should be aware of the water benefits to improve balance and core stability in disable populations.

References:

- Garcia, M. K., Joares, E. C., Silva, M. A., Bissolotti, R. R., Oliveira, S., & Battistella, L. R. (2012). The Halliwick Concept, inclusion and participation through aquatic functional activities. *Acta fisidtrica*, 19(3), 142–150.
- Tiroshi, R., Katz-Leurer, M., & Getz M. (2008). Halliwick-Based aquatic assessments: reliability and validity. *International Journal of Aquatic Research and Education*, 2, 224–236.

Effect of strength training in hypoxia on body composition and muscle performance

Ismael Martínez-Guardado¹, Marta Camacho-Cardenosa², Alba Camacho-Cardenosa¹, Guillermo Olcina¹, Rafael Timón¹

Sport Sciences Faculty. University of Extremadura. Caceres. wismi4@gmail.com

Resistance training is the best way to cause muscular adaptations. Current guidelines state that loads of $\geq 65\%$ 1RM are necessary to elicit favorable increases in hypertrophy (Kraemer et al., 2002; Kraemer & Ratamess, 2004). Besides this, the addition of hypoxic stimulus during resistance training is suggested to increase the metabolic responses, enhancing hypertrophy and muscle strength (Kon, Ikeda, Homma, & Suzuki, 2012). To determine the adaptations caused by hypoxia resistance training on strength and body composition. Sixteen untrained subjects participated in the study (weight: 74.68 ± 12.89 kg; height: 1.75 ± 0.08 cm; BMI: 24.28 ± 3.80 kg/m²). A training period of 7 weeks in a hypoxia chamber under normobaric hypoxia conditions ($FiO_2 = 13\%$) was performed. At the beginning, resistance training consisted of maximum repetitions to failure at 65% 1RM, and then, every two weeks, intensity training increased to finish at 80% 1RM. Initial and final measurements were taken. Body composition was determined through skin fold and muscle strength was evaluated with 1RM tests in the following exercises: bench press, biceps curl, french press, rowing hip and half squat. A general linear model of repeated measures was performed to observe the differences in every variable. Increases were observed in muscle percentage, so the fat percentage decreased significantly. They also showed an increase in the perimeters of the arm, leg and thigh with respect to their initial values. Regarding the performance values, subjects also increased the number of kg raised in 1RM for all exercises. The significance level was set at $p \leq 0.05$. In conclusion, resistance training of 7 weeks in hypoxia conditions increased strength and hypertrophy in untrained subjects.

Acknowledgement

This paper was supported by the Council of Extremadura (Aid for research groups GR 15020) for funding the research.

References:

- Kon, Ikeda, T., Homma, T., & Suzuki, Y. (2012). Effects of low-intensity resistance exercise under acute systematic hypoxia on hormonal responses. *Journal of Strength & Conditioning Research*, 26(6), 1–7.
- Kraemer, W. J., Adams, K., Cafarelli, E., Dudley, G. A., Dooly, C., Feigenbaum, M. S., Triplett-McBride, T. (2002). American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Medicine and Science in Sports and Exercise*, 34(2), 364–380.
- Kraemer, W. J., & Ratamess, N. A. (2004). Fundamentals of resistance training: progression and exercise prescription. *Medicine and Science in Sports and Exercise*, 36(4), 674–688.